

## NO SUBGROUP OF $A_4$ HAS INDEX 2

KEITH CONRAD

The group  $A_4$  has order 12, so its subgroups could have size 1, 2, 3, 4, 6, or 12. There are subgroups of orders 1, 2, 3, 4, and 12, but  $A_4$  has no subgroup of order 6 (equivalently, no subgroup of index 2). We will give three proofs that there is no subgroup of index 2, as corollaries of three different theorems from group theory.

**Theorem 1.** *If  $G$  is a finite group and  $N \triangleleft G$  then any element of  $G$  with order relatively prime to  $[G : N]$  lies in  $N$ . In particular, if  $N$  has index 2 then all elements of  $G$  with odd order lie in  $N$ .*

*Proof.* Let  $g$  be an element of  $G$  with order  $m$ , which is relatively prime to  $[G : N]$ . Reducing the equation  $g^m = e$  modulo  $N$  gives  $\bar{g}^m = \bar{e}$  in  $G/N$ . Also  $\bar{g}^{[G:N]} = \bar{e}$ , so the order of  $\bar{g}$  in  $G/N$  divides  $m$  and  $[G : N]$ . These numbers are relatively prime, so  $\bar{g} = \bar{e}$ , which means  $g \in N$ .  $\square$

**Corollary 2.** *There is no subgroup of index 2 in  $A_4$ .*

*Proof.* If  $A_4$  has a subgroup with index 2 then by Theorem 1, all elements of  $A_4$  with odd order are in the subgroup. But  $A_4$  contains 8 elements of order 3 (there are 8 different 3-cycles), and an index-2 subgroup of  $A_4$  has size 6, so not all elements of odd order can lie in the subgroup.  $\square$

**Theorem 3.** *If  $G$  is a finite group with a subgroup of index 2 then its commutator subgroup has even index.*

*Proof.* If  $[G : H] = 2$  then  $H \triangleleft G$ , so  $G/H$  is a group of size 2 and thus is abelian. So all commutators of  $G$  are in  $H$ , which means  $H$  contains the commutator subgroup of  $G$ . The index of the commutator subgroup therefore is divisible by  $[G : H] = 2$ .  $\square$

**Corollary 4.** *There is no subgroup of index 2 in  $A_4$ .*

*Proof.* We will show the commutator subgroup of  $A_4$  has odd index, so there is no index-2 subgroup by Theorem 3. The subgroup

$$V = \{(1), (12)(34), (13)(24), (14)(23)\}$$

is normal in  $A_4$  and  $A_4/V$  has size 3, hence is abelian, so the commutator subgroup of  $A_4$  is inside  $V$ . Each element of  $V$  is a commutator (e.g.,  $(12)(34) = [(123), (124)]$ ), so  $V$  is the commutator subgroup of  $A_4$ . It has index 3, which is odd.  $\square$

**Theorem 5.** *Every group of size 6 is cyclic or isomorphic to  $S_3$ .*

*Proof.* This is a special case of the classification of groups of order  $pq$  for primes  $p$  and  $q$ , but we give a self-contained treatment in this special case.

Let  $G$  have size 6 and assume  $G$  is not cyclic. We want to show  $G \cong S_3$ . By Cauchy,  $G$  contains elements  $a$  with order 2 and  $b$  with order 3. The subgroup  $H = \{1, a\}$  has index 3, so the usual left multiplication action of  $G$  on the left coset space  $G/H$  is a homomorphism

$G \rightarrow \text{Sym}(G/H) \cong S_3$ . If  $g$  is in the kernel then  $gH = H$ , so  $g \in H$ . Thus, if the kernel is nontrivial then it contains  $a$ . In particular,  $abH = bH$ . Since  $bH = \{b, ba\}$  and  $abH = \{ab, aba\}$ , either  $b = ab$  or  $b = aba$ . The first choice is impossible, so  $b = aba$ . Since  $a$  has order 2,  $ab = ba^{-1} = ba$ , which means  $a$  and  $b$  commute. Thus  $ab$  has order  $2 \cdot 3 = 6$ , so  $G$  is cyclic. We were assuming  $G$  is not cyclic, so the kernel of the map  $G \rightarrow \text{Sym}(G/H)$  is trivial, hence this is an isomorphism.  $\square$

**Corollary 6.** *There is no subgroup of index 2 in  $A_4$ .*

*Proof.* If  $A_4$  has an index-2 subgroup  $H$ , that subgroup has size 6 and therefore is isomorphic to either  $\mathbf{Z}/(6)$  or  $S_3$ . There are no elements in  $A_4$  with order 6, so the first choice is impossible:  $H$  must be isomorphic to  $S_3$ . In  $S_3$  there are three elements of order 2 (the transpositions). The group  $A_4$  also has only three elements of order 2  $((12)(34), (13)(24), (14)(23))$ , so these  $(2, 2)$ -cycles must lie in  $H$ . However, the elements of order 2 in  $S_3$  don't commute while the  $(2, 2)$ -cycles in  $A_4$  do commute, so we have a contradiction. Since  $H$  can't be isomorphic to  $S_3$ , it doesn't exist.  $\square$

For more proofs of this result, see [1].

#### REFERENCES

- [1] M. Brennan, D. Machale, Variations on a theme:  $A_4$  definitely has no subgroup of order six!, *Math. Mag.* **73** (2000), 36–40.